

# Nearshore Wave and Current Dynamics

Joan Oltman-Shay and Uday Putrevu  
Northwest Research Associates, Inc.  
14508 NE 20<sup>th</sup> Street  
Bellevue, WA 98007  
Phone: (425) 644 9660 Fax: (425) 644 8422  
Email: [joan@nwra.com](mailto:joan@nwra.com), [putrevu@nwra.com](mailto:putrevu@nwra.com)  
Award #: N0001496C0075

## LONG-TERM GOAL

The long-term goal of our research is to increase the understanding of nearshore (shoreline to nominally 15 m depth) fluid dynamics and to improve our predictive modeling of waves and currents in that region.

## SCIENTIFIC OBJECTIVES

In general terms, our work under this project has been geared towards contributing to the understanding and modeling of the large scale [ $O(10^2 - 10^3 \text{m}, 10^2 \text{s})$ ] fluid motions in the nearshore. Specifically, our recent work has concentrated on answering the following questions:

- 1) How does the alongshore variability of the bottom topography affect the longshore currents?
- 2) How important is the nonlinear interaction between edge waves?
- 3) What are the possible mechanisms that can explain the observations of shear waves on planar beaches?

## APPROACH

Our general approach is to use both field observations and theoretical/numerical models to increase our understanding of the large-scale fluid motions in the nearshore. In the work summarized below, we have primarily used theoretical models to answer the questions mentioned above. Our field program is part of another project; the work carried out under that project is being reported separately (see also the section titled "Related Projects" below). Data collected by our field program will be used in the near future to further our study of wave and current theory and predictive models.

## WORK COMPLETED

We have successfully completed the following:

- 1) Longshore currents – the effects of alongshore variation of topography: a) An evaluation of the limitations of using a simple model to evaluate these effects (Sancho *et al.*, 1998) and b) An extension of the dispersive mixing of momentum to include these effects and to remove the assumption of steady state (Putrevu and Svendsen, 1998).

2) Infragravity waves: a) An evaluation of how nonplanar topography affects edge-wave kinematics (dispersion relationship and spatial structure) (Putrevu and Oltman-Shay, 1998) and b) A general formulation of the equations governing the nonlinear interactions of edge waves (Kirby *et al.*, 1998).

3) Shear waves: a) An evaluation of the effects of forcing on shear wave generation (Haller *et al.*, 1998) and b) An evaluation of how lateral mixing changes the stability of longshore currents (Putrevu *et al.*, 1998).

## RESULTS

Sancho, Svendsen, and Putrevu, 1998 (Modeling of longshore currents over longshore nonuniform topographies: Effects of second-order terms, in revision, *J. Geophys Res.*). Putrevu *et al.* (1995) found that alongshore inhomogeneities of the bottom topography induce alongshore pressure gradients which can significantly influence the longshore currents. For the case in which the alongshore variations of the bottom topography are weak, Putrevu *et al.* suggested that the alongshore pressure gradient can be calculated in a simple way. The present work was designed to evaluate the limitations of the simple model proposed by Putrevu *et al.* The results show that the Putrevu *et al.* model works in cases in which the alongshore variations in the bottom topography vary over lengths that are long in comparison with the surf zone width. For cases in which the alongshore variation of the topography occurs over relatively short distances (like, *e.g.*, in a rip-channel) the simple model of Putrevu *et al.* does not work even if the absolute magnitude of the changes is relatively small.

Putrevu and Svendsen, 1998 (Shear dispersion in the nearshore, under review, *European Journal of Mechanics*). In this work, we extended the results of Svendsen and Putrevu (1994) to the general case in which the assumptions of alongshore uniformity and steady state are abandoned. This work showed that it is possible to account for the dispersive-mixing effects of the vertical nonuniformity of the short-wave-averaged velocity field over an arbitrary bottom topography without resorting to a fully three-dimensional calculation. The results, however, are far more complicated than the results for the simple situation considered by Svendsen and Putrevu (1994). In particular, this work shows that the results obtained by Svendsen and Putrevu (1994) represent only the leading term of the complete result. The importance of these additional terms is at present unknown.

Putrevu and Oltman-Shay, 1998 (Influence functions for edge wave propagation over a nonplanar bathymetry, *Physics of Fluids*, 10, 330-332). In this work we investigated how nonplanar features influence the propagation of edge waves. To do this, we assumed that the bottom topography over which the infragravity waves propagate was broken down into a base profile (*e.g.*, a planar profile) and a deviation from the base profile. The edge-wave quantities (*i.e.*, surface elevation and wavenumber) were similarly expanded. We found that when these expansions are substituted into the edge-wave equations, the lowest order edge-wave problem reduces to the problem of edge waves propagating over the base profile (whose solution is known by assumption), and the next order problem gives us the corrections to this base solution. We found that the correction to the dispersion relationship is proportional to the (cross-shore) integral of the product of the bottom perturbation and an "influence function". This influence function has its maximum at the shoreline and decays away from the shore. Also, the magnitude of the influence function increases with edge-wave mode. These results show that the dispersion relationship is more sensitive to the features at the shoreline and quite insensitive to features at moderate distances from the shore, thus explaining the differences between Holman and Bowen's (1979) and Kirby *et al.*'s (1981) results. Our results also lead us to conclude that the higher modes are more sensitive to shoreline features than the lower modes. The effect of deviations from planar topography for the spatial structure similarly can be expressed in terms of influence functions, but are slightly more complicated.

Kirby, Putrevu, and Özkan-Haller, 1998 (Evolution equations for edge waves and shear waves on longshore uniform beaches, *Intl. Conf. Coastal Engrg.*, Copenhagen) In this work, we derived the general equations that govern the interaction of edge and shear waves on longshore-uniform beaches. We showed that edge waves on open coastal beaches satisfy a complex array of triad resonance conditions, thereby providing a pathway for energy to move across a range of frequencies and mode numbers. Preliminary calculations on planar beaches (in the absence of longshore currents) indicate that the time scales for energy exchange between realistic edge wave modes can be as short as ten wave periods, suggesting that the mechanism is comparable in speed to the mechanisms causing edge wave growth. In addition, these calculations showed the following intriguing result: triads involving at least one opposite-going wave lead to significant transfers, while triads involving three waves all propagating in the same direction do not produce an interaction. We are presently investigating the reasons for this lack of interaction and are also extending our calculations to other topographies to determine whether the lack of interaction of three edge waves all propagating in the same direction is a general result.

Haller, Putrevu, Dalrymple, and Oltman-Shay, 1998 (Wave group forcing of low frequency surf zone motion, under revision, *J. Geophys. Res.*; Low-frequency surf zone response to wave groups, *Intl. Conf. Coastal Engrg.*, Copenhagen). In this work, we investigated the effects that wave group forcing has on shear instabilities in the nearshore region. This work was motivated by the results of Dodd *et al.* (1992) and Shrira *et al.* (1997). Dodd *et al.* found that Bowen and Holman's (1989) instability theory fails to explain the observations of shear waves on planar beaches. A related prediction is that on barred beaches there is a low-frequency, low-wavenumber cutoff below which no shear wave generation takes place. (The theory predicts that the instabilities will be damped out due to friction for realistic values of the bottom friction.) Shrira *et al.* showed that shear waves that are expected to be damped out by friction in the Bowen and Holman theory could grow to significant amplitudes due to triad interactions provided their initial amplitudes exceeded a certain critical value. However, how such initial (small amplitude) shear waves are generated remains unexplained in the work by Shrira *et al.* In this work we showed that the direct forcing by wave groups sets up oscillations that are remarkably similar in all respects to shear waves. We further showed that the forced response is extremely strong and could easily provide the initial amplitudes required by the Shrira *et al.* model. We then analyzed field data from the NSTS and SUPERDUCK experiments and showed that there is significant evidence that the spatial and temporal scales of forcing required to set up these initial oscillations does exist in the field.

Putrevu, Kirby, Oltman-Shay, and Özkan-Haller, 1998 (On the viscous destabilization of longshore currents, *Intl. Conf. Coastal Engrg.*, Copenhagen). This work was motivated by essentially the same concerns that motivated the work described in the previous paragraph. Here we investigated how the inclusion of lateral mixing changes the stability characteristics of longshore currents. We first extended Bowen and Holman's (1989) formulation to include lateral mixing. We then solved the resulting equations for the model problem considered by Bowen and Holman. This solution showed that the lateral mixing significantly alters the stability characteristics of the longshore current (see figure). In particular, it eliminates the low-frequency, low-wavenumber cutoff predicted by the inviscid theory. Thus, it is plausible that lateral mixing could be an important factor in explaining the observations of shear waves in cases where the inviscid calculations predict stability (and hence no shear wave generation).




Figure caption: Plot of growth rate vs. wavenumber for the model problem considered by Bowen and Holman (1989). The lines marked BH89 represent the solution given by Bowen and Holman (1989). Note that Bowen and Holman's solution predicts that there is zero growth for small wavenumbers, indicating that there is no shear wave generation at these wavenumbers. The lines marked "Present solution" represent the variation of growth rate when lateral mixing is included. Note that the inclusion of lateral mixing eliminates the low-frequency, low-wavenumber cutoff predicted by the inviscid theory.

## IMPACT/APPLICATION

Scientific results that will influence the modeling of nearshore waves and currents include the following:

It is important to account for alongshore nonuniformities of the bottom topography. Existing models of alongshore currents (which typically assume alongshore uniformity) easily can be extended to include minor alongshore variations as long as these variations occur over lengths that are much larger than the surf zone width.

It is possible to account for the dispersive mixing effects on the short-wave-averaged velocity field over arbitrary bottom topography without resorting to a fully three-dimensional calculation.

Either direct forcing or lateral mixing effects could potentially explain the observations of shear waves on planar beaches and could also account for the low-frequency, low-wavenumber shear waves observed on barred beaches.

The time scale of the nonlinear interactions between edge waves are comparable to the time scales over which edge waves grow due to direct forcing. Thus, any model that seeks to predict the edge wave climate in the nearshore (given the offshore wind-wave climate and the bottom bathymetry) should account for nonlinear interactions in addition to the direct forcing (due to modulations of the incident wind-waves).

## RELATED PROJECTS

1) Beach Probing System (BPS), funded by ONR Coastal Dynamics: The BPS measures the offshore (of the breakers) wind and infragravity wave field to estimate the inshore bathymetry and the wave and current conditions. This application is a direct result of basic research on nearshore infragravity waves conducted by the nearshore community of scientists under ONR Coastal Dynamics sponsorship. As part of this project, we just completed a large field experiment at Duck, NC and will be analyzing data from that experiment shortly. A report on this project is being submitted separately.

2) A field investigation of the dynamical importance of wind-wave groups in the surf zone, funded by NSF Physical Oceanography: We are collaborating with Prof. Ed Thornton (Naval Postgraduate School) and Prof. Ib Svendsen (University of Delaware) to analyze data collected during the 1997 SANDYDUCK experiment for evidence of vertical nonuniformity in the velocity field under infragravity waves due to forcing by wind-wave groupiness.

## REFERENCES

- Bowen, A. J., and R. A. Holman, 1989. Shear instabilities in the mean longshore current 1: Theory. *J. Geophys. Res.*, 94, 18,023-030.
- Dodd, N., J.M. Oltman-Shay and E.B. Thornton, 1992, "Shear instabilities in the longshore current: A comparison of observation and theory," *J. Phys. Oceanogr.*, 22, pp.62-82.
- Holman, R. A., and A. J. Bowen, 1979. Edge waves over complex beach profiles. *J. Geophys. Res.*, 84, 6330-6346.
- Kirby, J.T., R. A. Dalrymple and P. L.-F. Liu, 1981. Modification of edge waves by barred beach topography. *Coastal Engineering*, 5, 35-49.
- Putrevu, U., J. Oltman-Shay, and I. A. Svendsen, 1995. Effect of alongshore nonuniformities on longshore current predictions, *J. Geophys Res.*, 100, 16119-16130.
- Shrira, V. I., V. V. Vornovich, and N. G. Kozhelupova, 1997. Explosive instability of vorticity waves. *J. Phys. Oceanogr.*, 27, 542-554.
- Svendsen, I. A., and U. Putrevu, 1994. Nearshore mixing and dispersion. *Proc. Roy. Soc. A*, 445, 561-576.

## PUBLICATIONS

- Haller, M.C., U. Putrevu, R.A. Dalrymple, and J. Oltman-Shay, 1998 (under revision). Wave group forcing of low frequency surf zone motion, *J. Geophys. Res.*
- Haller, M.C., U. Putrevu, R.A. Dalrymple, and J. Oltman-Shay, 1998. Low-frequency surf zone response to wave groups, *Intl. Conf. Coastal Engrg.*, Copenhagen.
- Kirby, J.T., U. Putrevu, and H.T. Özkan-Haller, 1998. Evolution equations for edge waves and shear waves on longshore uniform beaches, *Intl. Conf. Coastal Engrg.*, Copenhagen

Putrevu, U., J.T. Kirby, J. Oltman-Shay, and H.T. Özkan-Haller, 1998. On the viscous destabilization of longshore currents, *Intl. Conf. Coastal Engrg.*, Copenhagen

Putrevu, U., and J. Oltman-Shay, 1998. Influence functions for edge wave propagation over a nonplanar bathymetry, *Physics of Fluids*, 10, 330-332.

Putrevu, U., and I.A. Svendsen, 1998 (under review). Shear dispersion in the nearshore, *European Journal of Mechanics*.

Sancho, F.E., I.A. Svendsen, and U. Putrevu, 1998 (under revision) Modeling of longshore currents over longshore nonuniform topographies: Effects of second-order terms, *J. Geophys Res.*